arXiv:1212.3241v1 [physics.chem-ph] 13 Dec 2012

Spin-charge and spin-orbital separations in density-functional theory

Daniel Vieira[1,*]

[1]*Departamento de Física, Universidade do Estado*
de Santa Catarina, Joinville, 89219-710 SC, Brazil

(Dated: December 14, 2012)

Abstract

It is known that the separation of electrons into spinons and chargons, the spin-charge separation, plays a decisive role when describing strongly correlated one-dimensional (1D) Friedel oscillations. Here, we extend the investigation by considering a third electron fractionalization: the separation into spinons and orbitons. Specifically, we deal with two exact constraints of exchange-correlation (XC) density-functionals: *(i)* The constancy of the highest occupied Kohn-Sham eigenvalues upon fractional electron numbers, and *(ii)* their discontinuities at integers. By means of 1D Hubbard chains, we show that spin-orbital separation can be decisive when dealing with derivative discontinuities of XC potentials, especially at strong correlations.

PACS numbers: 71.10.Fd, 71.15.Mb

I. INTRODUCTION

Spin and charge use to be treated as fundamental properties of ordinary electrons. However, as predicted in the 1960s, when confined in one dimension *interacting* electrons display the unusual property of separating their spin and charge into two independent quasiparticles: spinons and chargons.[1] Both behave just like ordinary electrons, but: spinons have spin-1/2 and no electrical charge, while chargons are spinless charged electrons. In the 1980s,[2,3] an additional fractionalization was shown to occur: the spin-orbital separation, for which spin and orbital degrees of freedom are decoupled to form the orbitons – particles with no spin and charge, carrying solely the orbital information. The spin-charge and spin-orbital separations have recent evidences of experimental observation,[3,4] teaching us that ordinary electrons can be considered bounded states of spinons, chargons and orbitons, three quasiparticles which can move independently and assume different velocities.

The Kohn-Sham (KS) formalism of density-functional theory (DFT),[5,6] by construction, retains the spin, charge and orbital degrees of freedom together, once it considers an auxiliary system of *noninteracting* particles. It has been shown that the separation into chargons and spinons are decisive when dealing with strongly correlated one-dimensional Friedel oscillations.[7] In this context, W. Yang *et. al.* have assumed strongly correlated systems as one of the modern challenges for DFT: *"The challenge of strongly correlated systems is a very important frontier for DFT. To fully understand the importance of these systems, they must be looked at within the wider realm of electronic structure methods."*[8] Here, we intend to give a contribution into this challenge: By means of the one-dimensional Hubbard model (1DHM),[9] we investigate the effect of electronic correlation on the spin-charge and spin-orbital separations. Specifically, we consider two exact constraints of exchange-correlation (XC) density-functionals: *(i)* the constancy of the highest occupied Kohn-Sham eigenvalues upon fractional electron numbers, and *(ii)* their discontinuities at integers. These constraints are usually not satisfied even by modern approaches, and are the cause of dramatic errors when dealing with any generic situation involving transport of charges. For example, the absence of a proper derivative discontinuity leads local and semi-local functionals to fail in yielding neutral atoms in molecular dissociation processes: instead, they predict dissociated subsystems with unphysical fractional charges.[10,11] This is usually associated to a difficulty in producing localized orbitals, resulting in excessive charge transfer which is sometimes referred to as delocalization error[12,13] of density functionals.

In detail, we shall compare the performance of the 1DHM local-density functional, its Perdew-Zunger self-interaction correction (PZSIC) and its spin-charge separation correction (SCSC). At strong correlations, we show that spin-orbital separation can be decisive when dealing with derivative discontinuities of XC potentials.

II. THEORETICAL BACKGROUND

(1) The one-dimensional Hubbard model: In one dimension, and in second-quantized notation, the Hubbard model[9] is defined as

$$\hat{H} \;=\; -t\sum_{j,\sigma}^{L}\left(c^{\dagger}_{j\sigma}c_{j+1,\sigma}+\text{H.c.}\right)+U\sum_{j}^{L}c^{\dagger}_{j\uparrow}c_{j\uparrow}c^{\dagger}_{j\downarrow}c_{j\downarrow}+\sum_{j,\sigma}^{L}V_{j}c^{\dagger}_{j\sigma}c_{j,\sigma}, \tag{1}$$

where L is the number of sites, t is the amplitude for hopping between neighboring sites and V_j is a local external potential acting on site j. Occupation of each site is limited to two particles, necessarily of opposite spin. For this reason, in spite of using the XC nomenclature, we shall deal only with correlation, especially in situations for which its major counterpart is due to the local (on-site) interaction U.

The basic Hohenberg-Kohn and KS theorems of DFT hold for the 1DHM too, once the density $n(\mathbf{r})$ is replaced by the on-site occupation $n(j)$.[14] In terms of this variable, local-(spin)-density approximations for Hubbard chains and rings have been constructed.[15–17] Here, we chose the BALDA/FN[17] as the reference functional, considering only non-polarized systems with $N_{\uparrow}=N_{\downarrow}$.

(2) Self-interaction correction: The BALDA/FN suffers from a self-interaction error. Perdew and Zunger have proposed the most known and used self-interaction correction (SIC),[18] written as:

$$E^{\text{PZSIC}}_{xc}[n]=E^{\text{BALDA}}_{xc}[n]\;-\;\sum_{\sigma=\uparrow,\downarrow}\sum_{k}^{N_\sigma}\left(E_H\left[n_{k\sigma}\right]+E^{\text{BALDA}}_{xc}\left[n_{k\sigma}\right]\right), \tag{2}$$

where $n_{k\sigma}(\mathbf{r}) \;=\; f_{k\sigma}|\psi_{k\sigma}(\mathbf{r})|^2$, for the KS orbitals $\psi_{k\sigma}$ and with occupation numbers $f_{k\sigma}$. The approach of Perdew and Zunger has a long history, including static[11,19,20] and time-dependent calculations.[21,22] Once the PZSIC functional of Eq. (2) depends explicitly on the KS orbitals, via the orbital densities $n_{k\sigma}$, we here shall employ the optimized effective potential (OEP) method[11] to obtain the orbital-independent PZSIC KS potential $v_{xc,\sigma}^{\mathrm{PZSIC}}[n](j)$ acting on each site j.

(3) Fractional electron numbers: In a system with $M = N + w$ $(0 \leq w \leq 1)$ electrons, the total ground-state energy is given by[23]

$$E(N + w) = (1 - w)E(N) + wE(N + 1). \tag{3}$$

Assuming that only the highest occupied (HO) KS orbital can be fractionally occupied, Janak[24] has proved that

$$\frac{\delta E}{\delta M} = \varepsilon_{\mathrm{HO}}(N) = \text{constant}, \tag{4}$$

where $\varepsilon_{k\sigma}$ are the KS eigenvalues. The derivative discontinuities of the energy at each integer N is then given by:

$$E_g(N) = IP(N) - EA(N) = \varepsilon_{N+1}(N + 1) - \varepsilon_N(N), \tag{5}$$

which is also referred to as fundamental energy gap.[25]

(4) The spin-charge separation correction: Considering spinons and chargons as independent entities, the occupied states will follow the semion distribution[26] of Fig. 1(b), where charge and spin were fractionalized to form the spin$-1/2$ spinons (as a sum of two $1/4$ spins) and the spinless chargons (as a sum of two $-e/2$ charges). As shown in Fig. 1(c), it has been proposed[7] that the occupied states of a non-interacting KS system are built by retaining spin and charge together, at expense of the presence of holons (the chargon antiparticles), whose densities are given by $\rho^+(j)$. The SCSC is then written as:

$$\begin{aligned} v_{\mathrm{KS}}^{\mathrm{SCSC}}[n](j) &= v_{ext}(j) + v_H[n](j) + v_{xc}^{\mathrm{BALDA}}[n](j) \\ &\qquad -v_H[\rho^+](j) - v_{xc}^{\mathrm{BALDA}}[\rho^+](j) \\ &\equiv v_{ext}(j) + v_H[n](j) + v_{xc}^{\mathrm{SCSC}}[n](j), \end{aligned} \tag{6}$$

with

$$\rho^+(j) = \sum_{\sigma=\uparrow,\downarrow} \frac{1}{2} \left[w|\psi_{N+1,\sigma}(j)|^2 + \sum_{k}^{N} |\psi_{k\sigma}(j)|^2 \right], \tag{7}$$


(a) Noninteracting system
(b) Interacting system
(c) Kohn-Sham system
Δ
spins + charges together
spinons + chargons + orbiton separated
spins + charges together + spinless holons + Δ
fractional occupation

FIG. 1: (Color online) Schematic representation of occupied orbitals for $4 \leq N \leq 5$. Fractional occupations are indicated by the dashed squares. Panel (b): The orbiton is indicated by the orbital excitation from level $k = 3$ to $k = 4$.

and

$$n(j) = w|\psi_{HO}(j)|^2 + \sum_{\sigma=\uparrow,\downarrow} \sum_{k}^{N_\sigma} |\psi_{k\sigma}(j)|^2, \tag{8}$$

where $0 \leq w \leq 1$, allowing fractional occupation.

III. RESULTS

Considering open Hubbard chains, in Figs. 2 (a)-(c) we plot $\varepsilon_{\mathrm{HO}}$ versus fractional charge occupations for three values of U. We note that BALDA and PZSIC yield a linear dependence for $\varepsilon_{\mathrm{HO}}$, indicating a parabolic behavior for the total energy functional. The slope of each straight line tend to increase as U is increased. In addition, while the exact (on Lanczos precision) energy gaps become larger as U is increased, the values of E_g yielded by LDA and PZSIC remain almost the same for all values of U, scaling to $\sim 50\%$ of the exact values when $U \gg 1.0t$. The SCSC, which includes the spin-charge separation mechanism and is also free from one-electron self-interaction error, performs much better than both, yielding almost constant values for $\varepsilon_{\mathrm{HO}}$, even though the values of E_g remain also constant when increasing U.

This difficulty in yielding accurate energy gaps is intrinsic of most available density functionals. In this context, based on the schematic representation of Figs. 1(b) and 1(c), we propose an alternative approach. Beyond spinons and chargons, in Fig. 1(b) we indicate an excitation from level $k = 3$ to $k = 4$. This is the schematic representation of an orbiton: an orbital excitation which behaves like a spinless and uncharged electron. When dealing with the noninteracting KS electrons of Fig. 1(c), the energy gaps should then be increased by a constant Δ, which is equivalent to include the presence of an "anti-orbiton", the extension of the SCSC idea of Eq. (6). Thus, in a system with spinons + chargons + orbitons, we propose the XC potential to be given by

$$v_{xc}^{approx+\Delta}[n](j) = v_{xc}^{approx}[n](j) + \Delta, \tag{9}$$

with

$$\Delta = \varepsilon_{\mathrm{HO}+\eta}^{\mathrm{KS}} - \varepsilon_{\mathrm{HO}}^{\mathrm{KS}}; \quad \eta = 0, 1, 2, ... \tag{10}$$

As conceptually thought, noninteracting systems should have $\eta = 0$, indicating the absence of spin-orbital separation. Strongly correlated systems, on the other hand, should have $\eta \geq 1$.

In Figs. 2 (d)-(f) we plot the results for the BALDA+Δ, PZSIC+Δ and SCSC+Δ approaches. When dealing with $U \lesssim 3t$, the best choice is $\eta = 0$. For stronger interactions, the values of E_g given by $\eta = 1$ are in great agreement with the exact values, multiplying the results of Figs. 2 (a)-(c) by a factor of ~ 2. Possibly, for $U \gg 10t$, it can be necessary to use $\eta \geq 2$. This is a clear delimitation between weakly and strongly correlated systems, whose connection between the increament of both, correlation and XC discontinuities, is in accordance with other well-known examples: In dissociation processes, to preserve neutrality of isolated atoms, the increment of atom-atom separation uses to be followed by a joint increment of electronic correlation and XC discontinuity.[8]

In Figs. 3 (a)-(c) we plot the XC potential profiles for $U = 5.0t$ and two slightly different electron numbers. When dealing with strong interactions, it has been shown that the XC potential should oscillate with $4k_F$.[7] Here, only SCSC yields the correct frequency, as seen by the four negative peaks in Fig. 3(c). The orbiton correction, as given by Eq. (9) with $\eta = 1$, induces a correct constant discontinuity in the XC potential when crossing $N = 4$, as seen by the jump from $N = 3.999$ to $N = 4.001$. The same is not observed for the XC potentials with $\eta = 0$, for which the curves are almost identical.

Density profiles: The density profiles of 1D Hubbard chains, which are invariant under the correction of Eq. (9), are known to suffer a $2k_F \to 4k_F$ crossover as U is increased.[7] Here, maintaining U fixed, we investigate the density profiles upon variation of electron numbers. In Figs. 3 (d)-(f) we show the density profiles for $2 \leq N \leq 4$, with an increase of 0.2 electrons for each curve, from bottom to top. The dashed lines indicate exact values (on Lanczos precision) for $N = 2$ and $N = 4$. The $4k_F$ Wigner crystal oscillations (identified by two or four peaks) are consequence of the strong interaction $U = 10.0t$. Only SCSC is able to reproduce the correct density profiles. Note that a similar change in frequency (from two to four peaks) could be obtained by fixing $N = 4$ and varying the interaction from $U = 0$ to $U \to \infty$.[7] Similar conclusions remain valid when including a confining harmonic potential (not showed here).

The emergence of $4k_F$ Wigner crystal oscillations, via Eq. (6), seems to be connected with the necessity of using Eq. (9), with $\eta \geq 1$. Both situations have a common point: The strongly interacting regime. W. Yang *el. al.* have recently pointed out[8] that, beyond strong correlations, it is a basic challenge to understand whether the KS orbitals and eigenvalues have any further significance. Therefore, the spinon-chargon-orbiton separation can be a first step into this direction.

In summary, specifically for the BALDA functional, we can conclude that: *(l)* spin-orbital separation, when included by considering the orbiton excitations, yields an increment of energy gaps; *(ll)* spin-charge separation, when included by means of the SCSC XC potential, yields $4k_F$ density oscillations and almost constant highest occupied KS eigenvalues. Considering *(l)* and *(ll)*, we can argue that when dealing with strong correlations, electrons should not be treated as unique particles. Instead, the separation into spinons, chargons and orbitons must be considered. The way to do so in a noninteracting KS calculation, which by construction retains spin, charge and orbital degrees of freedom together, is a combination of *(l)* and *(ll)*.

IV. CONCLUSIONS

Constancy of ε_{HO}*:* The BALDA and BALDA+PZSIC yielded typical results attributed to delocalization errors of density functionals: Incorrect linear behavior of $\varepsilon_{\mathrm{HO}}$ upon fractional charge occupation, with an increase in the slope when interaction is increased. On the



FIG. 2: (Color online) Highest occupied KS eigenvalues $\varepsilon_{\rm HO}$ obtained by means of Eq. (9). Panels (a)-(c): $\eta = 0$; Panels (d)-(f): $\eta = 1$.



FIG. 3: (Color online) (a)-(c): XC potential profiles for two slightly different electron numbers. (d)-(f): Density profiles for $2 \leq N \leq 4$. The dashed lines indicate exact values for $N = 2$ and $N = 4$, from bottom to top.

other hand, the BALDA+SCSC, which has been especially conceived to deal with strong interactions, yields almost constant values for ε_{HO}. The separation into spinons + chargons seems to be more efficient in correcting the BALDA than the Perdew-Zunger prescription.

Derivative discontinuity: Even though all functionals tested here yield accurate energy gaps at weak interactions, the accuracy is lost when interaction is increased. By means of Eq. (9), we proposed to also consider orbiton excitations. The results indicate a clear delimitation between weakly and strongly correlated systems: At strong correlations and with $\Delta \neq 0$, the inclusion of spin-charge and spin-orbital separations yields accurate energy gaps with correct $4k_F$ density profiles. It remains to be seen whether the same trends are observed in *ab initio* calculations under non-local $1/r$ Coulomb interactions. Understanding possible connections between using $\Delta \neq 0$ and other types of discontinuity corrections is a topic for further investigations.

Acknowledgments We thank Vivaldo Campo Jr. for his code of exact diagonalization and for the original version of the BALDA/FN code.

---

[*] Electronic address: danielv@joinville.udesc.br

[1] T. Giamarchi, *Quantum Physics in One Dimension* (Oxford University Press, Oxford, 2003).

[2] K. I. Kugel and D. I. Khomskii, Sov. Phys. Usp. **25**, 231 (1982).

[3] J. Schlappa, K. Wohlfeld, K. J. Zhou, M. Mourigal, M. W. Haverkort, V. N. Strocov, L. Hozoi, C. Monney, S. Nishimoto, S. Singh, A. Revcolevschi, J.-S. Caux, L. Patthey, H. M. Rnnow, J. van den Brink , and T. Schmitt, Nature **485**, 82 (2012).

[4] Y. Jompol, C. J. B. Ford, J. P. Griffiths, I. Farrer, G. A. C. Jones, D. Anderson, D. A. Ritchie, T. W. Silk, and A. J. Schofield, Science **325**, 597 (2009).

[5] W. Kohn, Rev. Mod. Phys. **71**, 1253 (1999).

[6] K. Capelle, Braz. J. Phys. **36**, 1318 (2006).

[7] D. Vieira, Phys. Rev. B **86**, 075132 (2012).

[8] A. J. Cohen, P. Mori-Sánchez, and W. Yang, Chem. Rev. **112**, 289 (2012).

[9] J. Hubbard, Proc. Roy. Soc. (London) A **276**, 238 (1963).

[10] A. Ruzsinszky, J. P. Perdew, G.I. Csonka, O. A. Vydrov, and G. E. Scuseria, J. Chem. Phys. **125**, 194112 (2006).

[11] S. Kümmel and L. Kronik, Rev. Mod. Phys. **80**, 3 (2008).

[12] A. J. Cohen, P. Mori-Sánchez, and W. Yang, Science **321**, 792 (2008).

[13] P. Mori-Sánchez, A. J. Cohen, and W. Yang, Phys. Rev. Lett. **100**, 146401 (2008).

[14] K. Schönhammer, O. Gunnarsson, and R. M. Noack, Phys. Rev. B **52**, 2504 (1995).

[15] N. A. Lima, M. F. Silva, L. N. Oliveira, and K. Capelle, Phys. Rev. Lett. **90**, 146402 (2003).

[16] V. V. França, D. Vieira and K. Capelle, New J. Phys. **14**, 073021 (2012).

[17] G. Xianlong, M. Polini, M. P. Tosi, V. L. Campo, K. Capelle, and M. Rigol, Phys. Rev. B **73**, 165120 (2006).

[18] J. P. Perdew and A. Zunger, Phys. Rev. B **23**, 5048 (1981).

[19] O. A. Vydrov, G.E. Scuseria, J.P. Perdew, A. Ruzsinszky, and G.I. Csonka, J. Chem. Phys. **124**, 094108 (2006).

[20] D. Vieira and K. Capelle, J. Chem. Theory Comput. **6**, 3319 (2010).

[21] D. Vieira, K. Capelle, and C. A. Ullrich, Phys. Chem. Chem. Phys. **11**, 4647 (2009).

[22] D. Hofmann, T. Körzdörfer, and S. Kümmel, Phys. Rev. Lett. **108**, 146401 (2012).

[23] J. P. Perdew, R. G. Parr, M. Levy, and J. L. Balduz, Jr., Phys. Rev. Lett. **49**, 1691 (1982).

[24] J. Janak, Phys. Rev. B **18**, 7165 (1978).

[25] K. Capelle, G. Vignale, and C. A. Ullrich, Phys. Rev. B **81**, 125114 (2010).

[26] A. Khare, *Fractional Statistics and Quantum Theory* (World Scientific Publishing, 2005).